



THE UNIVERSITY
OF KANSAS HOSPITAL
KUMED
3901 Rainbow Boulevard
Kansas City, Kansas 66160
PHYSICIAN'S ORDER FORM

Patient Label

TRANSESOPHAGEAL ECHO (TEE) ORDERS

Endocarditis prophylaxis is recommended prior to TEE for high risk patients only.

High Risk Category

- Prosthetic valves, including bioprosthesis and homograft valves
- Previous bacterial endocarditis
- Complex, cyanotic, congenital heart disease
- Surgically constructed systemic to pulmonary shunts

Prophylactic Regimens for Transesophageal Echocardiography

- | | |
|---|---|
| <ul style="list-style-type: none"> • Standard general prophylaxis • Unable to take oral medications • Allergic to penicillin | <p>Amoxicillin 2 g p.o. 1 h before procedure</p> <p>Ampicillin 2 g IV within 30 min before procedure</p> <p>Clindamycin 600 mg p.o. 1 h before procedure</p> <p>or</p> <p>Cephalexin 2 g p.o. 1 h before procedure</p> <p>or</p> <p>Azithromycin 500 mg p.o. 1 h before procedure</p> |
| <ul style="list-style-type: none"> • Allergic to penicillin and unable to take oral medications | <p>Clindamycin 600 mg IV within 30 min before procedure</p> <p>or</p> <p>Cefazolin 1 g IV within 30 min before procedure</p> |

TRANSESOPHAGEAL ECHO (TEE) ORDERS